

LEEDSTONE FOUNDATION 20 STARTER B60 DBZ6 MEDICATED

Type C Medicated Feed
GRAIN MIXTURE FOR DAIRY STARTER --3 DAYS TO 3 MOS IN
CONFINEMENT

For control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in cattle up to 800 lbs. For the prevention of house, stable, face and horn fly emergence from the manure of treated cattle.

ACTIVE DRUG INGREDIENTS

Lasalocid (as Lasalocid sodium)	60.00 g/ton
Diflubenzuron.....	6.00 g/ton

GUARANTEED ANALYSIS

Crude Protein (Min).....	20.00 %
Crude Fat (Min).....	2.40 %
Crude Fiber (Max).....	7.00 %
Acid Detergent Fiber (ADF) (Max)	9.00 %
Calcium (Ca) (Min).....	0.75 %
Calcium (Ca) (Max).....	1.25 %
Phosphorus (P) (Min).....	0.54 %
Selenium (Se) (Min).....	0.46 ppm
Vitamin A (Min).....	8695 IU/LB

INGREDIENTS

Grain Products, Plant Protein Products, Processed Grain By-Products, Molasses Products, Calcium Carbonate, Roughage Products, Salt, Vegetable Oil Refinery Lipid, Dicalcium Phosphate, Monocalcium Phosphate, Artificial Flavor, Lignin Sulfonate, Xanthan Gum, Propionic Acid (a Preservative), Thiamine Mononitrate, Acetic Acid, Benzoic Acid (A Preservative), Zinc Sulfate, Vitamin E Supplement, Selenium Yeast, Zinc Amino Acid Complex, Sodium Silico Aluminate, Dried *Aspergillus oryzae* Fermentation Product, Manganese Amino Acid Complex, Copper Amino Acid Complex, Cobalt Carbonate, Sorbitan Monostearate, Vegetable Oil, Propyl Gallate (a preservative), Cobalt Glucoheptonate, Dried *Aspergillus niger* Fermentation Product, Vitamin A Supplement, Manganese Sulfate, Ferrous Sulfate, Sodium Hydroxide, Vitamin D3 Supplement, Zinc Oxide, Ammonium Hydroxide, Basic Copper Chloride, Manganous Oxide, Sorbic Acid (a Preservative), Methylparaben (a Preservative), Ethylenediamine Dihydroiodide, Propylparaben (a Preservative), Potassium Iodide, Dried *Trichoderma longibrachiatum* Fermentation Product, Sodium Aluminosilicate, Sodium Molybdate, Maltodextrins, Dried *Bacillus subtilis* Fermentation Product, Sodium Selenite, Dried *Enterococcus faecium* Fermentation Product, Dried *Lactobacillus casei* Fermentation Product.

R0HL-WIL-D 2

DIRECTIONS

When used as directed, this product provides sufficient diflubenzuron larvicide to prevent the emergence of adult house flies, stable flies, horn flies and face flies from manure of treated

See Reverse Side For Directions for Use and Precautionary Statements

MANUFACTURED FOR

Leedstone

222 County Rd 173, Melrose, MN 56352
Net Weight 50 lb (22.67 kg)

4548018-506

LEEDSTONE FOUNDATION 20 STARTER B60 DBZ6

R0HL

cattle. Existing adult flies will not be affected. Start feeding before flies appear and continue use until cold weather marks the end of the fly season.

Introduction of this product after an adult fly infestation is established will require treatment of cattle and confinement areas with adulticides if immediate elimination of the adult fly population is desired.

Hand feed continuously to cattle weighing 215 pounds at the rate of 3.25 pounds per head per day to provide 1 mg lasalocid per 2.2 lbs body weight per day (up to a maximum of 360 mg per head per day) and to provide 0.10 mg Diflubenzuron per kg of body weight.

CAUTION

KEEP OUT OF REACH OF CHILDREN Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse. Wear long-sleeved shirt and long pants, socks and shoes and gloves. **ENVIRONMENTAL HAZARDS** This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **Storage and Disposal:** Do not contaminate water, food or feed by storage or disposal. **Storage:** Store pesticide in original container in a dry place. **Product Disposal:** Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility. **Container Disposal:** Completely empty bag. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Store in a dry, well-ventilated area protected from rodents and insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, performance loss or death. This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.

USE ONLY AS DIRECTED

WARNING

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.