



**durvet**

Keep out of reach of children.

For Oral Use In Horses Only

NDC 30798-837-81

**DURAMECTIN**<sup>TM</sup>  
(ivermectin) **PASTE 1.87%**

**Anthelmintic & Boticide**

**Net Wt. 0.21 oz (6.08 g)**

Removes worms and bots with a single dose.

Contents will treat up to 1250 lb body weight.

ANADA# 200-326, Approved by FDA

**Removes Worms & Bots**

**Removes Worms & Bots**



**durvet**

www.durvet.com

**DURAMECTIN**<sup>TM</sup>  
(ivermectin) **PASTE 1.87%**

**Anthelmintic & Boticide**



7 45801 11070 0

Manufactured by: Bimeda-UTC Animal Health Inc.  
Cambridge, Ontario, Canada N3C 2V4  
Distributed by: Durvet Inc.  
Blue Springs, MO 64014  
www.durvet.com  
Rev. 07.12 80UR020C

**INDICATIONS:** Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. DuraMectin<sup>TM</sup> (ivermectin) Paste provides effective treatment and control of the following parasites in horses. **Large Strongyles** (adults)- *Strongylus vulgaris* (also early forms in blood vessels), *S. edentatus* (also tissue stages), *S. equinus*, *Tricodontophorus* spp. including *T. brevicauda* and *T. serratus* and *Craterostomum acuticaudatum*; **Small Strongyles** (adults, including those resistant to some benzimidazole class compounds)- *Coronocylus* spp. including *C. coronatus*, *C. labiatus* and *C. labratus*, *Cyathostomum* spp. including *C. catinatum* and *C. pateratum*; *Cylicocylus* spp. including *C. insigne*, *C. leptostomum*, *C. nassatus*, and *C. brevicapsulatus*; *Cylicodontophorus* spp.; *Cylicostephanus* spp. including *C. calicatus*, *C. goldi*, *C. longibursatus*, and *C. minutus* and *Petrovinema poculatum*; **Small Strongyles**- Fourth-stage larvae; **Pinworms** (adults and fourth-stage larvae)- *Oxyuris equi*; **Ascarids** (adults, third-stage and fourth-stage larvae)- *Parascaris equorum*; **Hairworms** (adults)- *Trichostrongylus axei*; **Large-mouth Stomach Worms** (adults)- *Habronema muscae*; **Bots** (oral and gastric stages)- *Gasterophilus* spp. including *G. intestinalis* and *G. nasalis*; **Lungworms** (adults and fourth-stage larvae)- *Dictyocaulus arnfieldi*; **Intestinal Threadworms** (adults)- *Strongyloides westeri*; **Summer Sores** caused by *Habronema* and *Draschia* spp. cutaneous third-stage larvae; Dermatitis caused by **neck threadworm** microfilariae, *Onchocerca* sp.

**DOSAGE AND ADMINISTRATION:** This syringe contains sufficient paste to treat one 1250 lb horse at the recommended dose rate of 91 mcg ivermectin per lb (200 mcg/kg) body weight. Each weight marking on the syringe plunger delivers enough paste to treat 250 lb body weight.

(1) While holding plunger, turn the knurled ring on the plunger 1/4 turn to the left and slide it so the side nearest the barrel is at the prescribed weight marking. (2) Lock the ring in place by making a 1/4 turn to the right. (3) Make sure that the horse's mouth contains no feed. (4) Remove the cover from the tip of the syringe. (5) Insert the syringe tip into the horse's mouth at the space

between the teeth. (6) Depress the plunger as far as it will go, depositing paste on the back of the tongue. (7) Immediately raise the horse's head for a few seconds after dosing.

**PARASITE CONTROL PROGRAM:** All horses should be included in a regular parasite control program with particular attention being paid to mares, foals and yearlings. Foals should be treated initially at 6 to 8 weeks of age, and routine treatment repeated as appropriate. Consult your veterinarian for a control program to meet your specific needs. DuraMectin<sup>TM</sup> (ivermectin) Paste effectively controls gastrointestinal nematodes and bots of horses. Regular treatment will reduce the chances of verminous arteritis caused by *Strongylus vulgaris*.

**PRODUCT ADVANTAGES: Broad-spectrum Control** - DuraMectin<sup>TM</sup> (ivermectin) Paste kills important internal parasites, including bots and the arterial stages of *S. vulgaris*, with a single dose.

DuraMectin<sup>TM</sup> (ivermectin) Paste is a potent antiparasitic agent that is neither a benzimidazole nor an organophosphate.

**ANIMAL SAFETY:** DuraMectin<sup>TM</sup> (ivermectin) Paste may be used in horses of all ages, including mares at any stage of pregnancy. Stallions may be treated without adversely affecting their fertility.

**WARNING: Do not use in horses intended for human consumption.**

**Not for use in humans. Keep this and all drugs out of reach of children.** Refrain from smoking and eating when handling. Wash hands after use. Avoid contact with eyes. The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. To report adverse reactions in users, to obtain more information or to obtain MSDS, contact Durvet at 1-800-821-5570.

**PRECAUTIONS:** DuraMectin<sup>TM</sup> (ivermectin) Paste has been formulated specifically for use in horses only. This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.

**Environmental Safety:** Ivermectin and excreted ivermectin residues may adversely affect aquatic organisms. Do not contaminate ground or surface water. Dispose of the syringe in an approved landfill or by incineration.

**INFORMATION FOR HORSE OWNERS:** Swelling and itching reactions after treatment with DuraMectin<sup>TM</sup> (ivermectin) Paste have occurred in horses carrying heavy infections of neck threadworm (*Onchocerca* sp. microfilariae). These reactions were most likely

the result of microfilariae dying in large numbers. Symptomatic treatment may be advisable. Consult your veterinarian should any such reactions occur.

Healing of summer sores involving extensive tissue changes may require other appropriate therapy in conjunction with treatment with DuraMectin<sup>TM</sup> (ivermectin) Paste. Reinfection, and measures for its prevention, should also be considered. Consult your veterinarian if the condition does not improve.

**STORAGE INFORMATION: STORE BETWEEN 20°C - 25°C (68°F - 77°F). Excursions between 15°C - 30°C (59°F - 86°F) are permitted.**

TAKE TIME



OBSERVE LABEL  
DIRECTIONS

Anthelmintic & Boticide

For Oral Use In Horses Only

**DURAMECTIN**<sup>TM</sup>  
(ivermectin) **PASTE 1.87%**

For Oral Use In Horses Only

**DURAMECTIN**<sup>TM</sup>  
(ivermectin) **PASTE 1.87%**

Anthelmintic & Boticide